

**Hizentra**<sup>®</sup>  
Immune Globulin Subcutaneous  
(Human) 20% Liquid

Biotherapies for Life<sup>®</sup> **CSL Behring**

Hizentra is an Ig\* therapy that provides proven protection for PI with the convenience of self-administration, so you can focus on everyday living

\*Ig=Immunoglobulin



## My Life, My Way With Hizentra

**Participate from the comfort of your HOME in a LIVE webinar educational program for PI patients and their family members and/or caregivers**

This webinar educational program for **primary immune deficiency (PI)** patients and their family members and/or caregivers will feature a presentation from a healthcare professional who helps treat PI, stories from a PI patient just like you, and an opportunity to ask questions live.

**This webinar educational program offers PI patients and their family members and/or caregivers the opportunity to:**

- Learn from a healthcare professional about the self-administration of Hizentra
- Hear from a Patient Advocate<sup>†</sup> sharing his/her experience with PI and journey to Hizentra
- Gain valuable information and deepen their understanding of Hizentra through a **LIVE** Q&A session

### WEBINAR EDUCATIONAL PROGRAMS

**Wednesday, August 28, 2019**

**Wednesday, October 16, 2019**

**Wednesday, December 11, 2019**

7:00 PM – 8:00 PM ET

**Register now to reserve your spot!**

**[www.HizentraPatientProgram.com](http://www.HizentraPatientProgram.com)**

**1-866-480-4637**

### Nurse Speaker



**Carla Duff, ARNP, CPNP-PC, MSN, CCRP, IgCN**  
University of South Florida,  
Department of Pediatrics  
St. Petersburg, FL

### Patient Advocate Speaker



**Carl Schlacht**  
Cleveland, OH

<sup>†</sup>Patient Advocates are not healthcare professionals or medical experts. For medical questions, patients should contact their healthcare provider. Advocates are compensated by CSL Behring LLC for their time and/or expenses.

### Important Safety Information

**WARNING: Thrombosis (blood clots) can occur with immune globulin products, including Hizentra. Risk factors can include: advanced age, prolonged immobilization, a history of blood clotting or hyperviscosity (blood thickness), use of estrogens, installed vascular catheters, and cardiovascular risk factors.**

**If you are at high risk of blood clots, your doctor will prescribe Hizentra at the minimum dose and infusion rate practicable and will monitor for signs of clotting events and hyperviscosity. Always drink sufficient fluids before infusing Hizentra.**

**See your doctor for a full explanation, and the full prescribing information for complete boxed warning.**

Hizentra is a prescription medicine used to treat:

- Primary immune deficiency (PI) in patients 2 years and older
- Chronic inflammatory demyelinating polyneuropathy (CIDP) in adults

**Please see Important Safety Information on reverse and full Prescribing Information available at this presentation.**

## Important Safety Information (continued)

Treatment with Hizentra might not be possible if your doctor determines you have hyperprolinemia (too much proline in the blood), or are IgA-deficient with antibodies to IgA and a history of hypersensitivity. Tell your doctor if you have previously had a severe allergic reaction (including anaphylaxis) to the administration of human immune globulin. Tell your doctor right away or go to the emergency room if you have hives, trouble breathing, wheezing, dizziness, or fainting. These could be signs of a bad allergic reaction.

Inform your doctor of any medications you are taking, as well as any medical conditions you may have had, especially if you have a history of diseases related to the heart or blood vessels, or have been immobile for some time. Inform your physician if you are pregnant or nursing, or plan to become pregnant.

**Infuse Hizentra under your skin *only*; do not inject into a blood vessel.** Self-administer Hizentra only after having been taught to do so by your doctor or other healthcare professional, and having received dosing instructions for treating your condition.

Immediately report to your physician any of the following symptoms, which could be signs of serious adverse reactions to Hizentra:

- Reduced urination, sudden weight gain, or swelling in your legs (possible signs of a kidney problem).
- Pain and/or swelling or discoloration of an arm or leg, unexplained shortness of breath, chest pain or discomfort that worsens on deep breathing, unexplained rapid pulse, or numbness/weakness on one side of the body (possible signs of a blood clot).
- Bad headache with nausea; vomiting; stiff neck; fever; and sensitivity to light (possible signs of meningitis).
- Brown or red urine; rapid heart rate; yellowing of the skin or eyes; chest pains or breathing trouble; fever over 100°F (possible symptoms of other conditions that require prompt treatment).

Hizentra is made from human blood. The risk of transmission of infectious agents, including viruses and, theoretically, the Creutzfeldt-Jakob disease (CJD) agent and its variant (vCJD), cannot be completely eliminated.

The most common side effects in the clinical trials for Hizentra include redness, swelling, itching, and/or bruising at the infusion site; headache; chest, joint or back pain; diarrhea; tiredness; cough; rash; itching; fever, nausea, and vomiting. These are not the only side effects possible. Tell your doctor about any side effect that bothers you or does not go away.

Before receiving any vaccine, tell immunizing physician if you have had recent therapy with Hizentra, as effectiveness of the vaccine could be compromised.

**Please see full prescribing information for Hizentra, including boxed warning and the patient product information.**

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

Biotherapies for Life® **CSL Behring**

Hizentra is manufactured by CSL Behring AG and distributed by CSL Behring LLC.  
Hizentra® and MyHizentra® are registered trademarks of CSL Behring AG.  
Biotherapies for Life® is a registered trademark of CSL Behring LLC.

©2019 CSL Behring LLC. The product information presented on this invitation is intended for US residents only. HIZ-0739-FEB19

*Anyone under 18 years of age must be accompanied by a parent or caregiver.*

  
**Hizentra®**  
Immune Globulin Subcutaneous  
(Human) 20% Liquid